

Operator-algebraic approach to the yrast spectrum of weakly interacting trapped bosons

Masahito Ueda¹ and Tatsuya Nakajima^{2,*}

¹ *Department of Physics, Tokyo Institute of Technology, Meguro-ku, Tokyo 152-8551, Japan, and CREST, Japan Science and Technology Corporation (JST), Saitama 332-0012, Japan*

² *Department of Physics, The University of Texas, Austin, Texas 78712, USA*

(August 16, 2001)

We present an operator-algebraic approach to deriving the low-lying quasi-degenerate spectrum of weakly interacting trapped N bosons with total angular momentum $\hbar L$ for the case of $L/N \ll 1$, demonstrating that the lowest-lying excitation spectrum is given by $27gn_3(n_3 - 1)/34$, where g is the strength of the repulsive contact interaction and n_3 the number of excited octupole quanta. Our method provides constraints for these quasi-degenerate many-body states and gives higher excitation energies that depend linearly on N .

I. INTRODUCTION

Realization of Bose-Einstein condensates (BECs) in atomic gases [1–3] has stimulated explosively growing interest in several subfields of physics. With unprecedented controllability in creating, manipulating, and probing systems, many previously “gedanken” experiments have become or are expected to become a reality [4]. Another characteristic feature of gaseous BEC systems is their good isolation from the surrounding environment, as the system is suspended in a vacuum chamber by a magnetic or an optical trapping potential. When the potential is axisymmetric, the projected angular momentum (AM) on the symmetry axis is conserved and can take on any integral multiple of $\hbar$. In contrast, in bulk superfluid systems, the container plays the role of a highly dissipative environment so that the system can sustain AM only at thermodynamically stable values, resulting in the quantization of circulation. In this respect, gaseous BEC systems are more like nuclear systems than bulk superfluid systems, and it is of interest to study the energy spectrum of a weakly interacting BEC system as a function of AM, namely the yrast spectrum [5–8].

The lowest-energy state of an isolated system subject to a given AM, $\hbar L$, is called the yrast state [9]. When the system is in the yrast state, all kinetic energy is used for rigid-body rotation, so the system is at zero temperature. This allows one to measure fine energy levels—the yrast spectrum—close to the ground state. Recently, the yrast state has been extensively discussed in the context of gaseous BEC systems [5,8,10–16]. Mottelson has pointed out that for repulsive contact interactions the yrast spectrum is dominated by quasi-degenerate quadrupole and octupole excitations [8], and Kavoulakis *et al* have shown that the interaction between octupole excitations makes the excitation energy of this mode slightly higher than that of the quadrupole mode [10]. Bertsch and Papenbrock [12] have found numerically that for $0 \leq L \leq N$, the energy of the yrast state is given by

$E = L\hbar\omega + gN(N - 1 - L/2)$, where ω is the frequency of the confining potential, g the strength of the interaction, and N the number of bosons, which we shall assume to be constant. The corresponding eigenstate has been shown to exist in Refs. [13–15]. Recently, we have found numerically that the *lowest-lying* yrast spectrum arises from the pairwise repulsive interaction between octupole excitations and that the corresponding interaction energy is given by $1.59gn_3(n_3 - 1)/2$, where n_3 is the number of octupole excitations [16]. In this paper, we present an operator-algebraic approach [17] to analytically deriving these results in a systematic way, demonstrating that the interaction energy between octupole excitations is given by $27gn_3(n_3 - 1)/34$, where the coefficient $27/34$ agrees precisely with the numerically obtained value [16]. Our method also provides constraints for the quasi-degenerate many-body states and gives higher excitation energies that depend linearly on N .

This paper is organized as follows. Section II describes the model under consideration and formulates the problem. Section III describes collective modes and discusses why they behave as well-defined quasiparticles when the system undergoes the Bose-Einstein condensation. Section IV derives the yrast spectrum based on our operator-algebraic approach, and Sec. V concludes this paper.

II. FORMULATION OF THE PROBLEM

We consider a two-dimensional system of harmonically trapped N bosons interacting via a contact δ -function interaction. The Hamiltonian of the system is given by

$$H = \sum_{i=1}^N \left[-\frac{\hbar^2}{2M} \left(\frac{\partial^2}{\partial x_i^2} + \frac{\partial^2}{\partial y_i^2} \right) + \frac{M\omega^2}{2}(x_i^2 + y_i^2) \right] + \frac{2\pi\hbar g}{M\omega} \sum_{i \neq j} \delta^{(2)}(\mathbf{r}_i - \mathbf{r}_j), \quad (1)$$

where M is the mass of the boson, ω the frequency of the confining potential, and g the strength of the interaction. In the second-quantized form, Eq. (1) is written as

$$\hat{H} = \int d^2z \left[\hat{\Psi}^\dagger(z) h(z) \hat{\Psi}(z) + \frac{2\pi\hbar g}{M\omega} \hat{\Psi}^\dagger(z) \hat{\Psi}^\dagger(z) \hat{\Psi}(z) \hat{\Psi}(z) \right], \quad (2)$$

where $z \equiv x + iy$ and

$$h(z) \equiv -\frac{2\hbar^2}{M} \frac{\partial^2}{\partial z \partial z^*} + \frac{M\omega^2}{2} |z|^2 \quad (3)$$

denotes the single-particle Hamiltonian.

Throughout this paper we consider the case of very weak interactions, *i.e.*, $|g|N \ll \hbar\omega$. It is then convenient to take the eigenfunctions of $h(z)$ as a basis set for expansion of the field operator. We introduce two sets of creation and annihilation operators,

$$\hat{a}_+ = \frac{z^*}{2d} + d \frac{\partial}{\partial z}, \quad \hat{a}_+^\dagger = \frac{z}{2d} - d \frac{\partial}{\partial z^*}, \quad (4)$$

$$\hat{a}_- = \frac{z}{2d} + d \frac{\partial}{\partial z^*}, \quad \hat{a}_-^\dagger = \frac{z^*}{2d} - d \frac{\partial}{\partial z}, \quad (5)$$

where $d \equiv \sqrt{\hbar/M\omega}$ is a characteristic length scale of the ground-state wave function of $h(z)$, and $\hat{a}_+$ (or $\hat{a}_-$) annihilates a quantum with magnetic quantum number 1 (or -1) and satisfies the canonical commutation relations

$$[\hat{a}_+, \hat{a}_+^\dagger] = [\hat{a}_-, \hat{a}_-^\dagger] = 1, \quad [\hat{a}_+, \hat{a}_-] = [\hat{a}_+, \hat{a}_-^\dagger] = 0. \quad (6)$$

In terms of these operators, h is diagonalized as

$$h = \hbar\omega(\hat{a}_+^\dagger \hat{a}_+ + \hat{a}_-^\dagger \hat{a}_- + 1). \quad (7)$$

The vacuum state $|0\rangle$ is defined by $\hat{a}_+|0\rangle = \hat{a}_-|0\rangle = 0$.

The operator corresponding to the magnetic quantum number m is given by $\hat{a}_+^\dagger \hat{a}_+ - \hat{a}_-^\dagger \hat{a}_-$. Because of the symmetry between $\hat{a}_+$ and $\hat{a}_-$, we may consider only the case of $m \geq 0$ with no loss of generality. Moreover, when $\hbar\omega \gg |g|N$, we may consider only those states $\{|\Psi\rangle\}$ that satisfy

$$\hat{a}_-^\dagger \hat{a}_- |\Psi\rangle = 0 \quad (8)$$

because for a given total AM, states that do not satisfy Eq. (8) are at least $2\hbar\omega$ higher in energy than those satisfying the equation. Physically, Eq. (8) suggests that no particle rotates in a clockwise direction. In what follows, we shall restrict ourselves to the Hilbert subspace constrained by Eq. (8), which corresponds to the lowest-Landau-level approximation in the theory of the fractional quantum Hall effect.

Within this approximation, a complete set of basis functions can be constructed as follows. Using $\langle z|\hat{a}_+|0\rangle = 0$ and Eq. (4), we obtain

$$\phi_0(z) \equiv \langle z|0\rangle = \frac{1}{\sqrt{\pi}d} e^{-|z|^2/2d^2}, \quad (9)$$

where the prefactor is determined so as to satisfy the normalization condition $\int d^2z |\phi_0(z)|^2 = 1$. The wavefunction $\phi_m(z)$ for the state with magnetic quantum number m is given by

$$\begin{aligned} \phi_m(z) &\equiv \langle z|m\rangle = \langle z| \frac{(\hat{a}_+^\dagger)^m}{\sqrt{m!}} |0\rangle = \frac{1}{\sqrt{m!}} \left(\frac{z}{2d} - d \frac{\partial}{\partial z^*} \right)^m \langle z|0\rangle \\ &= \frac{z^m}{\sqrt{\pi m!} d^{m+1}} e^{-|z|^2/2d^2}, \end{aligned} \quad (10)$$

and satisfies the orthonormal conditions

$$\int d^2z \phi_m^*(z) \phi_n(z) = \delta_{mn}. \quad (11)$$

It follows from Eq. (10) that $|z\rangle$ is a coherent state defined by

$$|z\rangle = (\sqrt{\pi}d)^{-1} \exp[(z^* \hat{a}_+^\dagger - z \hat{a}_+)/d] |0\rangle. \quad (12)$$

Expanding the field operator in terms of $\phi_m(z)$ as

$$\hat{\Psi}(z) = \sum_m \hat{b}_m \phi_m(z), \quad (13)$$

and substituting this into Eq. (2), we obtain

$$\hat{H} = \hbar\omega \hat{L} + g \sum_{m_1, \dots, m_4} V_{m_1 m_2 m_3 m_4} \hat{b}_{m_1}^\dagger \hat{b}_{m_2}^\dagger \hat{b}_{m_3} \hat{b}_{m_4}, \quad (14)$$

where

$$\hat{L} \equiv \sum_m m \hat{b}_m^\dagger \hat{b}_m \quad (15)$$

gives the total magnetic quantum number and $V_{m_1 m_2 m_3 m_4}$ is the matrix element of the interaction given as [12]

$$V_{m_1 m_2 m_3 m_4} = \frac{\delta_{m_1+m_2, m_3+m_4} (m_1+m_2)!}{2^{m_1+m_2} \sqrt{m_1! m_2! m_3! m_4!}}. \quad (16)$$

In Eq. (14) the term $\hbar\omega \hat{N}$ is dropped because the total number of bosons

$$\hat{N} \equiv \sum_m \hat{b}_m^\dagger \hat{b}_m \quad (17)$$

is conserved in our system.

Because $\hat{L}$ is also conserved in our system, we shall henceforth focus on the second term on the right-hand side (rhs) of Eq. (14), which will be denoted as $\hat{V} = g\hat{U}$. Our primary goal is to diagonalize $\hat{U}$ in operator form within the Hilbert subspace of non-negative m and under the two constraints

$$\hat{L} = L, \quad \hat{N} = N. \quad (18)$$

III. COLLECTIVE MODES

Mottelson has pointed out that low-lying excitations from the yrast line can be described in terms of collective modes excited by $(d^{-\lambda}/\sqrt{N\lambda!})\sum_{j=1}^N z_j^\lambda$, where λ is a positive integer. In the second-quantized form, the creation operators for the collective modes are written as

$$\begin{aligned}\hat{Q}_\lambda^\dagger &= \int d^2z \hat{\Psi}^\dagger(z) \frac{d^{-\lambda}}{\sqrt{N\lambda!}} z^\lambda \hat{\Psi}(z) \\ &= \frac{1}{\sqrt{N}} \sum_{m=0}^{\infty} \sqrt{\lambda+m} C_\lambda \hat{b}_{m+\lambda}^\dagger \hat{b}_m,\end{aligned}\quad (19)$$

where ${}_m C_n = m!/n!(m-n)!$. The commutation relation between $\hat{Q}_\lambda$ and the interaction Hamiltonian $\hat{V}$ is calculated to be

$$\begin{aligned}[\hat{V}, \hat{Q}_\lambda] &= \frac{2g}{\sqrt{N\lambda!}} \sum_{m_1, \dots, m_4} \frac{\delta_{m_1+m_2, m_3+m_4-\lambda} (m_1+m_2)!}{2^{m_1+m_2+\lambda} \sqrt{m_1! m_2! m_3! m_4!}} \\ &\times \left[2^\lambda \frac{m_4!}{(m_4-\lambda)!} - \frac{(m_1+m_2+\lambda)!}{(m_1+m_2)!} \right] \\ &\times \hat{b}_{m_1}^\dagger \hat{b}_{m_2}^\dagger \hat{b}_{m_3} \hat{b}_{m_4}.\end{aligned}\quad (20)$$

It follows from this relation that except for $\lambda = 1$, $\hat{Q}_\lambda$ does not commute with $\hat{V}$. The collective mode excited by $\hat{Q}_\lambda$ ($\lambda > 1$) is therefore not, in general, a well-defined quasiparticle. The only exception is the dipole operator $\hat{Q}_1$, for which the rhs of Eq. (20) vanishes. The dipole mode is therefore a well-defined collective mode, and for any L , an eigenstate of the Hamiltonian (14) may be constructed as [5,8]

$$|\Psi^L\rangle = (\hat{Q}_1^\dagger)^L |L=0\rangle, \quad (21)$$

where $|L=0\rangle$ denotes the exact ground state of the Hamiltonian (14) with zero total AM, and hence $\hat{V}|L=0\rangle = gN(N-1)|L=0\rangle$. The corresponding eigenenergy is given by

$$E^{\text{dipole}} = \hbar\omega L + gN(N-1) \quad (22)$$

because

$$\begin{aligned}\hat{H}|\Psi^L\rangle &= (\hbar\omega\hat{L} + \hat{V})(\hat{Q}_1^\dagger)^L |L=0\rangle \\ &= \hbar\omega L |\Psi^L\rangle + (\hat{Q}_1^\dagger)^L \hat{V} |L=0\rangle = E^{\text{dipole}} |\Psi^L\rangle.\end{aligned}$$

The state $|\Psi^L\rangle$ is the lowest-energy state of the Hamiltonian with AM L for the case of an attractive interaction ($g < 0$) and the highest-energy state for the case of a repulsive interaction ($g > 0$).

It should be noted that while $\hat{Q}_\lambda$ and $\hat{Q}_\mu$ commute for any λ and μ , $\hat{Q}_\lambda$ and $\hat{Q}_\mu^\dagger$ do not. In fact,

$$\begin{aligned}[\hat{Q}_\lambda, \hat{Q}_\mu^\dagger] &= \frac{1}{N} \sum_{m=0}^{\infty} \left[\theta(\lambda-\mu) \sqrt{m+\lambda} C_\lambda \sqrt{m+\lambda} C_\mu \hat{b}_m^\dagger \hat{b}_{m+\lambda-\mu} \right. \\ &\quad \left. + \theta(\mu-\lambda) \sqrt{m+\mu} C_\lambda \sqrt{m+\mu} C_\mu \hat{b}_{m+\mu-\lambda}^\dagger \hat{b}_m \right. \\ &\quad \left. - \sqrt{m+\lambda} C_\lambda \sqrt{m+\mu} C_\mu \hat{b}_{m+\mu}^\dagger \hat{b}_{m+\lambda} \right],\end{aligned}\quad (23)$$

where $\theta(x)$ is the Heaviside step function, which takes on 0, 1/2 and 1 for $x < 0$, $x = 0$ and $x > 0$, respectively. The commutation relation (23) implies that the collective modes carrying different angular momenta do not, in general, behave independently. For $\mu = \lambda$, Eq. (23) reduces to

$$\begin{aligned}[\hat{Q}_\lambda, \hat{Q}_\lambda^\dagger] &= \frac{1}{N} \sum_{m=0}^{\infty} ({}_{m+\lambda} C_\lambda - {}_m C_\lambda) \hat{b}_m^\dagger \hat{b}_m \\ &= \begin{cases} 1 & \text{for } \lambda = 1; \\ 1 + \frac{2L}{N} & \text{for } \lambda = 2; \\ 1 + \frac{3L}{2N} + \frac{3}{2N} \sum_m m^2 \hat{b}_m^\dagger \hat{b}_m & \text{for } \lambda = 3, \end{cases}\end{aligned}\quad (24)$$

where constraints (18) are used in deriving the second equality. The commutator $[\hat{Q}_\lambda, \hat{Q}_\lambda^\dagger]$ in Eq. (24) gives constants for $\lambda = 1, 2$, indicating that the dipole ($\lambda = 1$) and quadrupole ($\lambda = 2$) modes behave like bosons. In contrast, the commutator for the octupole ($\lambda = 3$) mode does not give a constant, indicating the anharmonicity of this mode. This anharmonicity leads to the interaction between octupole excitations, as shown in Sec. IV.

The energy minimization of $\hat{V}$ must be performed subject to constraints (18). In the case of $L/N \ll 1$, which we shall consider below, these constraints require that

$$\hat{n}_0 = \mathcal{O}(N), \quad \hat{n}_m = \mathcal{O}(N^0) \quad (m \geq 1), \quad (25)$$

where $\hat{n}_m \equiv \hat{b}_m^\dagger \hat{b}_m$ and $\mathcal{O}(N^k)$ denotes terms of the order of or less than N^k . In this case, $\hat{Q}_\lambda$ may be simplified as

$$\hat{Q}_\lambda \simeq \frac{1}{\sqrt{N}} \hat{b}_0^\dagger \hat{b}_\lambda, \quad (26)$$

and the commutation relations in Eq. (23) reduce to

$$[\hat{Q}_\lambda, \hat{Q}_\mu^\dagger] = \delta_{\lambda\mu} + \mathcal{O}\left(\frac{L}{N}\right). \quad (27)$$

The collective modes, therefore, behave as almost independent bosons for $L/N \ll 1$, with small inter-mode interactions on the order of L/N . These features are reflected explicitly in the yrast spectrum, as shown in the next section.

IV. YRAST SPECTRUM

Our basic idea for the derivation of the yrast spectrum is to rewrite the interaction Hamiltonian $\hat{V} = g\hat{U}$ under constraints (18) so that the excitation spectrum manifests itself. We first note that $\hat{U}$ can be rewritten as

$$\hat{U} = \sum_{m=0}^{\infty} \hat{A}_m^\dagger \hat{A}_m, \quad (28)$$

where

$$\hat{A}_m = 2^{-\frac{m}{2}} \sum_{k=0}^m \sqrt{m C_k} \hat{b}_k \hat{b}_{m-k} \quad (29)$$

may be regarded as the annihilation operator of a boson pair with the total magnetic quantum number m . As we shall consider only the case of $g > 0$, the energy minimization for $\hat{U}$ may be exploited to estimate the order of magnitude for each term.

The conditions (25) allow us to expand $\hat{A}_m$ as

$$\hat{A}_0 = \hat{b}_0^2, \quad \hat{A}_m = 2^{1-\frac{m}{2}} \hat{b}_0 \hat{b}_m + \mathcal{O}(N^0) \quad (m \geq 1). \quad (30)$$

Substituting these expansions into Eq. (28), we obtain

$$\hat{U} = N(N-1) - 2N \sum_{m=1}^{\infty} (1 - 2^{1-m}) \hat{n}_m + \mathcal{O}(N^{1/2}) \quad (31)$$

where the constraint $\hat{N} = N$ is used to eliminate $\hat{n}_0$. Because $\hat{n}_2$ and $\hat{n}_3$ appear on the rhs of Eq. (31) in a form proportional to $2\hat{n}_2 + 3\hat{n}_3$, the constraint $\hat{L} = L$ may be used to eliminate both of these terms simultaneously. We thus obtain

$$\begin{aligned} \hat{U} = & N \left(N - 1 - \frac{L}{2} \right) \\ & + N \left[\frac{\hat{n}_1}{2} + \sum_{m=4}^{\infty} \left(\frac{m}{2} - 2 + 2^{2-m} \right) \hat{n}_m \right] + \mathcal{O}(N^{1/2}). \end{aligned} \quad (32)$$

The minimum of $\hat{U}$ in Eq. (32) is attained when

$$\hat{n}_1 = \mathcal{O}(N^{-1/2}), \quad \hat{n}_m = \mathcal{O}(N^{-1/2}) \quad (m \geq 4). \quad (33)$$

These conditions are compatible with the constraint $\hat{N} = N$ because $\hat{n}_0 = \mathcal{O}(N^0)$, and with the other constraint $\hat{L} = L$ because it can be satisfied by an appropriate choice of $\hat{n}_2$ and $\hat{n}_3$. We note that the first term in Eq. (32) gives the yrast line [12].

Because the yrast line is dominated by the $m = 2$ and $m = 3$ modes, the low-lying excitations from the line due to binary interactions (14) should involve modes with at least up to $m = 6$. We therefore use $\hat{A}_m$ up to $m = 6$ without approximation:

$$\begin{aligned} \hat{A}_0 &= \hat{b}_0^2, \quad \hat{A}_1 = \sqrt{2} \hat{b}_0 \hat{b}_1, \quad \hat{A}_2 = \hat{b}_0 \hat{b}_2 + \frac{\hat{b}_1^2}{\sqrt{2}}, \\ \hat{A}_3 &= \frac{\hat{b}_0 \hat{b}_3}{\sqrt{2}} + \sqrt{\frac{3}{2}} \hat{b}_1 \hat{b}_2, \quad \hat{A}_4 = \frac{\hat{b}_0 \hat{b}_4}{2} + \hat{b}_1 \hat{b}_3 + \sqrt{\frac{3}{8}} \hat{b}_2^2, \\ \hat{A}_5 &= \frac{\hat{b}_0 \hat{b}_5}{\sqrt{8}} + \sqrt{\frac{5}{8}} \hat{b}_1 \hat{b}_4 + \frac{\sqrt{5}}{2} \hat{b}_2 \hat{b}_3, \\ \hat{A}_6 &= \frac{\hat{b}_0 \hat{b}_6}{4} + \sqrt{\frac{3}{8}} \hat{b}_1 \hat{b}_5 + \frac{\sqrt{15}}{4} \hat{b}_2 \hat{b}_4 + \frac{\sqrt{5}}{4} \hat{b}_3^2. \end{aligned} \quad (34)$$

Substituting these expressions into Eq. (28) yields

$$\begin{aligned} \hat{U} = & N \left(N - 1 - \frac{L}{2} \right) + \frac{27}{34} \hat{n}_3 (\hat{n}_3 - 1) + \hat{R} + \hat{S} \\ & + \frac{N}{2} \hat{B}_1^\dagger \hat{B}_1 + \frac{N}{4} \hat{B}_4^\dagger \hat{B}_4 + \frac{5N}{8} \hat{B}_5^\dagger \hat{B}_5 + \frac{17N}{16} \hat{B}_6^\dagger \hat{B}_6. \end{aligned} \quad (35)$$

Here $\hat{R}$ is given by

$$\begin{aligned} \hat{R} = & \sum_{k=4}^{\infty} \left(\frac{k-2}{2} \hat{n}_k^2 + \hat{n}_k \right) + \sum_{k=6}^{\infty} \frac{k-3}{2} \hat{n}_1 \hat{n}_k + \sum_{k=5}^{\infty} \frac{k-2}{2} \hat{n}_2 \hat{n}_k \\ & + \sum_{k=3}^{\infty} \sum_{l=k+1}^{\infty} \frac{k+l-4}{2} \hat{n}_k \hat{n}_l - 2(\hat{n}_3 + 1) \hat{n}_4 - \frac{5}{2} (\hat{n}_4 + 1) \hat{n}_5 \\ & - \frac{1}{2} (\sqrt{10} \hat{b}_3^\dagger \hat{b}_4^\dagger \hat{b}_2 \hat{b}_5 + 2\sqrt{5} \hat{b}_4^{\dagger 2} \hat{b}_3 \hat{b}_5 + \text{h.c.}) \\ & + \frac{1}{17} [23 \hat{n}_1 \hat{n}_5 + 32 \hat{n}_2 \hat{n}_4 \\ & - (14\sqrt{10} \hat{b}_1^\dagger \hat{b}_5^\dagger \hat{b}_2 \hat{b}_4 - \sqrt{30} \hat{b}_1^\dagger \hat{b}_5^\dagger \hat{b}_3^2 + 12\sqrt{3} \hat{b}_2^\dagger \hat{b}_4^\dagger \hat{b}_3^2 + \text{h.c.})], \end{aligned} \quad (36)$$

where h.c. denotes Hermitian conjugates of the preceding terms, $\hat{S}$ is given by

$$\hat{S} = \sum_{m=7}^{\infty} \left(\hat{A}_m^\dagger \hat{A}_m + \frac{m-4}{2} \hat{n}_0 \hat{n}_m \right), \quad (37)$$

and $\hat{B}_m$ ($m = 1, 4, 5, 6$) are defined by

$$\begin{aligned} \hat{B}_1 &\equiv \frac{1}{\sqrt{N}} \sum_{k=1}^5 \sqrt{k} \hat{b}_{k-1}^\dagger \hat{b}_k, \\ \hat{B}_4 &\equiv \frac{1}{\sqrt{N}} \left(\hat{b}_0 \hat{b}_4 - 2\hat{b}_1 \hat{b}_3 + \sqrt{\frac{3}{2}} \hat{b}_2^2 \right), \\ \hat{B}_5 &\equiv \frac{1}{\sqrt{N}} \left(\hat{b}_0 \hat{b}_5 - \frac{3}{\sqrt{5}} \hat{b}_1 \hat{b}_4 + \sqrt{\frac{2}{5}} \hat{b}_2 \hat{b}_3 \right), \\ \hat{B}_6 &\equiv \frac{1}{\sqrt{N}} \left(\hat{b}_0 \hat{b}_6 + \frac{\sqrt{6}}{17} \hat{b}_1 \hat{b}_5 + \frac{\sqrt{15}}{17} \hat{b}_2 \hat{b}_4 + \frac{\sqrt{5}}{17} \hat{b}_3^2 \right). \end{aligned} \quad (38)$$

Note that no approximation is made in obtaining Eq. (35). It follows from (33) that $\hat{R} = \mathcal{O}(N^{-1/2})$. The operator $\hat{S}$ is positive semidefinite. To minimize it, we consider the states that satisfy

$$\hat{b}_k |\Psi\rangle = o(N^{-1/2}) \quad (k \geq 7), \quad (39)$$

so as to give $\langle \Psi | \hat{S} | \Psi \rangle = o(N^0)$, where $o(N^\alpha)$ denotes terms whose order of magnitude is less than N^α . Under the same condition we obtain

$$\begin{aligned} \hat{U} = & N \left(N - 1 - \frac{L}{2} \right) + \frac{27}{34} \hat{n}_3 (\hat{n}_3 - 1) \\ & + \frac{N}{2} \hat{B}_1^\dagger \hat{B}_1 + \frac{N}{4} \hat{B}_4^\dagger \hat{B}_4 + \frac{5N}{8} \hat{B}_5^\dagger \hat{B}_5 \\ & + \frac{17N}{16} \hat{B}_6^\dagger \hat{B}_6 + o(N^0), \end{aligned} \quad (40)$$

which constitutes the primary result of our paper.

Let us define the *quasi-degenerate* states, $\{|\Psi\rangle\}$, as the ones satisfying both condition (39) and

$$\hat{B}_m|\Psi\rangle = o(N^{-\frac{1}{2}}) \quad (m = 1, 4, 5, 6). \quad (41)$$

The following constraints on these states are then obtained:

$$\langle \hat{B}_m^\dagger \hat{B}_m \rangle = o(N^{-1}) \quad (m = 1, 4, 5, 6), \quad (42)$$

$$\langle \hat{n}_m \rangle = o(N^{-1}) \quad (m \geq 7). \quad (43)$$

The expectation values of $\hat{U}$ for these states are therefore given by

$$\langle \hat{U} \rangle = N \left(N - 1 - \frac{L}{2} \right) + \frac{27}{34} n_3(n_3 - 1) + o(N^0), \quad (44)$$

where n_3 is the number of octupole excitations, as $\hat{Q}_\lambda^\dagger \hat{Q}_\lambda = \hat{n}_\lambda + \mathcal{O}(N^{-1/2})$ for $\lambda \geq 1$. Since the population of the octupole mode with $n_3 \geq 2$ raises the energy, while that of the quadrupole mode does not [10,16], the ground-state configuration is given by $n_2 = L/2$ and $n_3 = 0$ when L is even and by $n_2 = (L-3)/2$ and $n_3 = 1$ when L is odd. These results are valid up to the order of N^0 , as Eq. (44) is valid up to the same order.

As seen in Eq. (40), the quasi-degenerate octupole spectrum lies above the yrast states. Although the octupole mode couples with other modes through

$$\begin{aligned} [\hat{b}_3, \hat{U}] - \frac{27}{17} \hat{b}_3^\dagger \hat{b}_3^2 &= \frac{\sqrt{3N}}{2} \hat{b}_2 \hat{B}_1 + \sqrt{N} \hat{B}_1^\dagger \hat{b}_4 - \frac{\sqrt{N}}{2} \hat{b}_1^\dagger \hat{B}_4 \\ &\quad + \frac{\sqrt{10N}}{8} \hat{b}_2^\dagger \hat{B}_5 + \frac{\sqrt{5N}}{8} \hat{b}_3^\dagger \hat{B}_6, \end{aligned} \quad (45)$$

the expectation value of the rhs over the quasi-degenerate state satisfying Eq. (41) is on the order of $o(N^0)$. The quasi-degenerate spectrum $27n_3(n_3 - 1)/34$ is therefore valid up to the same order.

The representation of $\hat{U}$ in Eq. (40) not only describes the quasi-degenerate yrast spectra, as discussed above, but also explains higher excitations whose energies are linear in N . Since

$$[\hat{B}_m, \hat{B}_n] = \mathcal{O}(N^{-1/2}) \quad \text{and} \quad [\hat{B}_m, \hat{B}_n^\dagger] = \delta_{mn} + \mathcal{O}(N^{-1/2}), \quad (46)$$

each $\hat{B}_m$ ($m = 1, 4, 5, 6$) behaves as a boson-like operator and satisfies

$$\hat{B}_m^\dagger \hat{B}_m = \hat{n}_m + \mathcal{O}(N^{-1/2}). \quad (47)$$

Because of the definitions (38) of $\hat{B}_m$ and from (33), the excitation energies for $\hat{B}_4$, $\hat{B}_5$, and $\hat{B}_6$ are expected to be

$$\begin{aligned} \epsilon_0 + \epsilon_4 - 2\epsilon_2 &= gN/4, \\ \epsilon_0 + \epsilon_5 - \epsilon_2 - \epsilon_3 &= 5gN/8, \quad \text{and} \\ \epsilon_0 + \epsilon_6 - 2\epsilon_3 &= 17gN/16, \end{aligned} \quad (48)$$

respectively, and the excitation energies for $\hat{B}_1$ are expected to be

$$\epsilon_1 + \epsilon_2 - \epsilon_0 - \epsilon_3 = gN/2 \quad \text{and} \quad 2\epsilon_1 - \epsilon_0 - \epsilon_2 = gN \quad (49)$$

for $n_1 = 1$ and $n_1 = 2$, respectively. Here

$$\epsilon_\lambda = -2gN(1 - 2^{1-\lambda}) \quad (50)$$

gives the energy of a single excitation of $\hat{Q}_\lambda$ ($\lambda \geq 1$) [8] and $\epsilon_0 \equiv 0$. In fact, these excitation energies appear as the coefficients for $\hat{B}_m^\dagger \hat{B}_m$ in Eq. (40).

Here we discuss the physical meaning of Eq. (41). In the BEC the $m = 0$ state is macroscopically occupied. When the BEC has angular momentum L ($1 \ll L \ll N$), the occupation numbers of $m = 2$ and $m = 3$ states become of the order of L . Because of the interactions, the following scattering processes, whose energy contributions are of the order of $\mathcal{O}(N^0)$, become important: $0+2 \leftrightarrow 1+1$, $0+3 \leftrightarrow 1+2$, $0+4 \leftrightarrow 2+2$, $0+5 \leftrightarrow 2+3$, and $0+6 \leftrightarrow 3+3$. The corresponding terms in the interaction Hamiltonian can be diagonalized by the introduction of the quasiparticle operators B_m ($m = 1, 4, 5, 6$) defined in Eq. (38). This procedure is nothing but the renormalization of the bare operators b_m due to the interaction. The true vacuum is therefore not the bare one defined by $b_m|\Psi\rangle = 0$ ($m = 1, 4, 5, 6$) but that of the quasiparticles defined by $B_m|\Psi\rangle = 0$. This is the physical meaning of the constraint (41).

On the other hand, the contact interaction dictates that neither quadrupole ($m = 2$) nor octupole ($m = 3$) mode can make $\mathcal{O}(N^0)$ energy contributions together with higher angular momentum states ($m = 7$ or higher). Hence the supplementary constraint (39) for $m \geq 7$ should be imposed for the complete definition of the (interacting) vacuum state. In fact, from Eq. (37) b_m excitations are shown to need energy costs which are linear in N . We note that because of this renormalization procedure of the quasiparticle operators, our excitation spectrum achieves the accuracy of $o(N^0)$ for quasi-degenerate states. Moreover it should be noted that the other low-lying excitations can also be explained naturally in terms of B_m and b_m excitations.

As an independent check of the above analytical results, we have numerically performed an exact diagonalization of the Hamiltonian, confirming that the constraints (43) on the quasi-degenerate states are satisfied. The excitations with energies linear in N are also found to exist and are shown in Fig. 1 as open circles. Here the solid line indicates the yrast line $-gNL/2$, and the dashed, dotted, dash-dotted, and dash-double-dotted lines correspond, respectively, to the $gN/4$, $gN/2$, $5gN/8$, and $17gN/16$ excitations that are predicted by Eq. (40). The broken curve connects the points of collective excitations $\hat{Q}_\lambda^\dagger|0\rangle$. We see that the excitations shown in Fig. 1 can be well explained in terms of single excitations of $\hat{B}_m$ ($m = 1, 4, 5, 6$). When these higher modes are excited, the quasi-degenerate conditions (41) are no longer

met, so that the coupling of these modes to the octupole mode through Eq. (45) becomes significant. This coupling causes $\mathcal{O}(N^0)$ -order corrections to the excitation energies, which are on the order of $\mathcal{O}(N)$. The examples of our numerical calculations demonstrating these small corrections are shown in Table I. Since the relative order of magnitude of these corrections is on the order of $1/N$, they cannot be discerned in Fig. 1.

Our method provides some constraints on the quasi-degenerate states, including the yrast state, through (39) and (41). From these constraints, we may gain some insight into quantum fluctuations occurring in the quasi-degenerate states $\{|\Psi\rangle\}$. For example, from the condition $\hat{B}_1|\Psi\rangle = o(N^{-\frac{1}{2}})$ and (33), we obtain

$$\left(\hat{b}_0^\dagger \hat{b}_1 + \sqrt{2} \hat{b}_1^\dagger \hat{b}_2 + \sqrt{3} \hat{b}_2^\dagger \hat{b}_3\right) |\Psi\rangle = o(N^0). \quad (51)$$

This equation strongly suggests that the interconversions between the states with $m = 0, 1, 2, 3$ (and the interconversions between quadrupole and octupole modes [16]) play an important role in the quantum fluctuations of the quasi-degenerate states. We stress here that the rhs of Eq. (51) is not $\mathcal{O}(N^0)$ but $o(N^0)$. Similarly, from the condition $\hat{B}_5|\Psi\rangle = o(N^{-\frac{1}{2}})$, we obtain

$$\left(\hat{b}_0 \hat{b}_5 + \frac{\sqrt{10}}{5} \hat{b}_2 \hat{b}_3\right) |\Psi\rangle = o(N^0). \quad (52)$$

It follows from these two constraints that on a mean-field level we have $\langle \hat{n}_1 \rangle \simeq 15 \langle \hat{n}_5 \rangle / 2$. From the constraints $\hat{B}_m|\Psi\rangle = o(N^{-\frac{1}{2}})$ ($m = 4, 5, 6$), we may also have $\langle \hat{n}_4 \rangle \simeq 3 \langle \hat{n}_2 \rangle^2 / 2N$, $\langle \hat{n}_5 \rangle \simeq 2 \langle \hat{n}_2 \rangle \langle \hat{n}_3 \rangle / 5N$, and $\langle \hat{n}_6 \rangle \simeq 5 \langle \hat{n}_3 \rangle^2 / 289N$, respectively. Except for the numerical coefficient in the last relation, these results agree with the mean-field analysis in Ref. [10], although these approximate relations are not always satisfied by exact numerical results.

V. CONCLUSIONS

In this paper we have proposed an operator-algebraic approach to analyzing the yrast spectrum of weakly interacting bosons. The basic idea is to exploit the macroscopic occupation of a particular mode to rewrite the second-quantized interaction Hamiltonian under the restriction of the total particle-number conservation and the angular-momentum conservation.

We have analytically shown that the lowest-lying excitation spectrum from the yrast line is given by $27gn_3(n_3 - 1)/34$. Equation (40) also shows that this quasi-degenerate energy spectrum is separated from other levels by energy gaps that are linear in N . These results are in agreement with what we have found by numerical diagonalization of the same Hamiltonian [16]. Equation (40) can also explain the excitations whose energies are linear in N , again in agreement with our exact diagonalization calculations [16].




FIG. 1. Energies (in units of gN) of the yrast states and those of low-lying excited states with $N = 10000$ for $1 \leq L \leq 8$. Open circles show our numerical results, the solid line indicates the yrast line $-gNL/2$, and the broken curve connects the collective modes excited by $\hat{Q}_\lambda^\dagger$. The dashed, dotted, dash-dotted, and dash-double-dotted lines correspond, respectively, to the $gN/4$, $gN/2$, $5gN/8$, and $17gN/16$ excitations predicted by Eq. (40).

The method presented herein only exploits the presence of a macroscopically occupied state, and it should therefore not be restricted to the case of $L/N \ll 1$ discussed above. Such an extension is in progress and will be reported elsewhere.

ACKNOWLEDGMENTS

M.U. acknowledges support by a Grant-in-Aid for Scientific Research (Grant No. 11216204) by the Ministry of Education, Science, Sports, and Culture of Japan, and by the Toray Science Foundation. T.N. was supported for research in the USA by the Japan Society for the Promotion of Science.

Note added: After completion of this work, we became aware of three studies (cond-mat/0011303, cond-mat/00011430, and cond-mat/00012438) that used different methods to derive the same analytic formula $27gn_3(n_3 - 1)/34$, as presented herein.

* Permanent address: Physics Department, Graduate School of Science, Tohoku University, Sendai 980-8578, Japan.

- [1] M.H. Anderson, J.R. Ensher, M.R. Mathews, C.E. Wieman, and E.A. Cornell, *Science* **269**, 198 (1995).
- [2] K.B. Davis, M.-O. Mewes, M.R. Andrews, N.J. van Druten, D.S. Durfee, D.M. Kurn, and W. Ketterle, *Phys. Rev. Lett.* **75**, 3969 (1995).
- [3] C.C. Bradley, C.A. Sackett, J.J. Tollett, and R.G. Hulet, *Phys. Rev. Lett.* **75**, 1687 (1995); *ibid.* **79**, 1170 (1997) (E); C.C. Bradley, C.A. Sackett, and R.G. Hulet, *ibid.* **78**, 985 (1997).
- [4] F. Dalfovo, S. Giorgini, L.P. Pitaevskii and S. Stringari, *Rev. Mod. Phys.* **71**, 463 (1999); for a collection of review articles, see “Bose-Einstein Condensation in Atomic Gases”, edited by M. Inguscio, S. Stringari and C. Wieman (IOS Press, Amsterdam, 1999).
- [5] N.K. Wilkin, J.M.F. Gunn and R.A. Smith, *Phys. Rev. Lett.* **80**, 2265 (1998).
- [6] D.A. Butts and D.S. Rokhsar, *Nature* **397**, 327 (1999).
- [7] M. Linn and A.L. Fetter, *Phys. Rev. A* **60**, 4910 (1999); D.L. Feder, C.W. Clark and B.I. Schneider, *ibid.* **61**, 11601 (2000); *Phys. Rev. Lett.* **82**, 4956 (1999).
- [8] B. Mottelson, *Phys. Rev. Lett.* **83**, 2695 (1999).
- [9] I. Hamamoto, in *Treatise on Heavy-Ion Science*, edited by D. A. Bromley (Plenum, New York, 1985), vol.3, p.313; I. Hamamoto and B. Mottelson, *Nucl. Phys.* **A507**, 65 (1990).
- [10] G.M. Kavoulakis, B. Mottelson and C.J. Pethick, *Phys. Rev. A* **62**, 63605 (2000).
- [11] A.D. Jackson, G.M. Kavoulakis, B. Mottelson and S.M. Reimann, *Phys. Rev. Lett.* **86**, 945 (2001).
- [12] G.F. Bertsch and T. Papenbrock, *Phys. Rev. Lett.* **83**, 5412 (1999).
- [13] A.D. Jackson and G.M. Kavoulakis, *Phys. Rev. Lett.* **85**, 2854 (2000).
- [14] R.A. Smith and N.K. Wilkin, *Phys. Rev. A* **62**, 61602 (2000).
- [15] T. Papenbrock and G.F. Bertsch, *Phys. Rev. A* **63**, 23616 (2001).
- [16] T. Nakajima and M. Ueda, *Phys. Rev. A* **63**, 43610 (2001).
- [17] This analytic result was first presented by us at the symposium on “Physics and applications of laser cooling” of the annual meeting of the Physical Society of Japan (Niigata, Sep. 25, 2000).

L	7	7	7	8	8	8	8
excitation energy	2502.8	5000.0	6253.3	2507.7	5000.0	6254.0	10628

TABLE I. Excitation energies measured from the yrast line with $N = 10000$ for $L = 7$ and 8. The numerical results agree with our analytical predictions of $N/4 = 2500$, $N/2 = 5000$, $5N/8 = 6250$, and $17N/16 = 10625$ with $\mathcal{O}(N^0)$ deviations that arise from the coupling to the octupole mode through Eq. (45).